

Evaluation of a modified GnRH-based TAI protocol associated with estrus detection in cyclic beef heifers inseminated with sex-selected semen



M.G. Colazo¹, P. Whittaker², D. Bignell¹, and R.J. Mapletoft³

¹Livestock Research Section, Alberta Agriculture and Forestry, Edmonton, Canada. ²Leduc Veterinary Hospital, Leduc, Canada. ³WCVM, University of Saskatchewan, Saskatoon, Canada.


BACKGROUND & OBJECTIVE

- The initial gonadotropin-releasing hormone (GnRH) treatment in a 5-day Co-synch protocol for timed-AI (TAI) has been shown to be unnecessary in cyclic dairy and beef heifers inseminated with conventional semen (Theriogenology 2011;76:578; Reprod Fertil Dev 2015;28:135).
- This study compared pregnancy per AI (P/AI) between sex-selected and conventional semen in cyclic beef heifers subjected to the modified (without initial GnRH) 5-day Co-synch protocol and evaluated the usefulness of an estrus detection (ED) patch to identify heifers that were most likely to conceive.

MATERIALS & METHODS




- 1034 beef heifers (325 to 523 kg of body weight and 13 to 15 months of age).
- Three locations, 2 sires, sex-selected or conventional semen, and 1 technician.
- Ultrasonography to determine cyclicity and reproductive normalcy (Day 0) and pregnancy status (27 or 48 d after TAI).

Figure 1. Illustration of treatments.



CIDR= insert containing 1.38 g P4 (CIDR, Zoetis Animal Health)
 PGF_{2α} = 500 µg cloprostenol i.m. (Estroplan, Vetoquinol NA Inc.)
 GnRH= 100 µg i.m. (Fertiline, Vetoquinol NA Inc.)
 ED patch = Estrotec™ (Estrotec Inc.)

Table 1. Estrus detection patches scoring based on color change between application (Day 5) and TAI (Day 8).

Score 0	Score 1	Score 2	Score 3
			
Unchanged	< 50% change	≥ 50% change	Missing

- Statistical analyses: PROC GLIMMIX in SAS 9.3.

RESULTS

- 890 cyclic and normal heifers were used; CIDR retention rate was 98.0%.
- P/AI did not differ (P>0.05) between sires (58.4 vs. 58.0%), but it was greater (P<0.01) for conventional semen than sex-selected semen (63.7 vs. 52.6%).
- Overall, P/AI was greater (P<0.01) when ED patches were scored 2 (64.8%) than 0 (42.9%) or 1 (46.8%), with score 3 (missing) intermediate (58.5%).
- P/AI with sex-selected semen was greater (P<0.05) when ED patches were scored 2 or 3 (Table 2).

Table 2. Effect of ED patch score and type of semen on P/AI.

	Score 0 (n=170)	Score 1 (n=79)	Score 2 (n=526)	Score 3 (n=65)
Sex-selected, %	36.1 ^a	37.2 ^a	59.8 ^b	58.3 ^b
Conventional, %	49.4 ^c	58.3 ^{cd}	69.6 ^d	58.6 ^{cd}

^{a,b}Within a row, percentage without a common superscript differed (P<0.05).
^{c,d}Within a row, percentage without a common superscript differed (P<0.01).

SUMMARY

- P/AI was greater in heifers inseminated with conventional semen.
- The Estrotec ED patches could be used to identify animals for TAI with sex-selected semen.

This study was supported by Alberta Agriculture and Forestry, Vetoquinol NA Inc., Estrotec Inc. and collaborative beef producers.